- **References**:-

1. Krell R. Value-Added Products from Beekeeping; FAO Agricultural Service Bulletin No 124; Food and Agriculture Organization of the United Nations: Rome, Italy, (**1996),** Available online: http://www.fao.org/docrep/w0076E/w0076E00.htm.

2.CiuluM., Nadia S., Maria I. P. and Gavino S. **(2016)**, Recent Advances in the Analysis of phenolic compounds in unifloral honeys.*Molecules*, 21: (451), 1 - 32.

3.StrayerS. E, Karen E. and Shaun K. **(2014)**,*Food Protection Trends* 8-14.

4. RaoC. R. M, Lakkineni C. A. K., and Chandra B. S. **(2015)**, Quantitative Analysis of Oxytetracycline Residues in Honey by High Performance Liquid Chromatography.,*International Research Journal of Biological Sciences* Vol. 4(5), 59-65.

5.KivrakI, ŞeydaK, and Mansur H.**(2016)**,Development of a rapid method for the determination of antibiotic residues in honey using UPLC-ESI-MS/MS, *Food Science and Technol.,* 36(1): 90-96.

6.Pena A, PelantovaN, LinoC. M.,Silveira M. I. N. and SolichP. **(2005)**,Validation of an Analytical Methodology for Determination of Oxytetracycline and Tetracycline Residues in Honey by HPLC with Fluorescence Detection,*J. Agric. Food Chem.*,53, 3784-3788.

7.Johnson S, Jadon N, Mathur H. B. and Agarwal H. C.**(2010)**,Antibiotic residues in honey, *Centre for Science and Environment*, New Delhi, 01-48.

8.GangulyN. K. and Chandy S. J. **(2011)**, Rationalizing antibiotic use to limit antibiotic resistance in India,*Indian J Med Res.* 134(3): 281–294.

9.Fuselli S.R., Gende L. B., Garcia S. B., Eguaras M. J. and Fritz R. **(2005),** Inhibition of *Paenibacillus Larvae* Subsp*Larvae b*y the essential oils of two wild plants and their emulsifying agents, *Span. J. Agri. Res.* **3** (2):220-224.

10.Tillotson G. S, Doern G.V. and Blondeau J. M. (2006),Optimal antimicrobial therapy: the balance of potency and exposure, *Exp. Opi. Invest. Drugs* **15**, 335-337.

11.Al-WailiN,Khelod S., Ahmed A.and Mohammad J. A. **(2012)**, Antibiotic, pesticide, and microbial contaminants of honey: Human Health Hazards; *The ScientificWorld Journal*, doi:10.1100/2012/930849.

12.GumustasM,KurbanogluS,Bengi U,Sibel A. O.**(2013)**, UPLC versus HPLC on drug analysis: advantageous applications and their validation parameters,*Chromatographia.,* 76, 1365–1427.

13.Tylova T,Novak J. O., Miroslav F.**(2010)**, High-throughput analysis of tetracycline antibiotics and their epimers in liquid hog manure using Ultra Performance Liquid Chromatography with UV detection,*Chemosphere*, 78:353–359.

14. PaillerJ.Y,Krein A,Pfister L, Hoffmann L. and GuignardC.**(2009)**, Solid phase extraction coupled to liquid chromatography-tandem mass spectrometry analysis of sulfonamides, tetracyclines, analgesics and hormones in surface water and wastewater in Luxembourg; *Science of the Total Enviro.*, 407:4736–4743.

15. Thompson T. S, Noot D. K, Cavert J. and Pernal S. F. **(2003)**, Determination of lincomycin and tylosin residues in honey using solid phase extraction and liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. J. Chromatogr. A., 1020, 241-250.

16. Sheridan R, Policastro B, Thomas S. and Rice D. (2008), Analysis and occurrence of 14 sulfonamide antibacterials and chloramphenicol in honey by solid-phase extraction followed by LC/MS/MS analysis. *J. Agric. Food Chem.*, 56(10), 3509-3516.

17. TamosiunasV. and Padarauskas A.**(2008)**, Comparison of LC and UPLC Coupled to MS–MS for the Determination of Sulfonamides in Egg and Honey,*Chromatographia* 67,783-788.

18. JinY, Zhang J, Wen Z,Wenwen Z, Lin W, Jinhui Z. and Yi L.**(2017)**, Development and validation of a multiclass method for the quantification of veterinary drug residues in honey and royal jelly by liquid chromatography–tandem mass spectrometry, *Food Chemistry* 221, 1298–1307.

19.GuidiL. R., Luiza H. M, Silva C. F., Nicki J. and Engeseth M. B. A. **(2015)**, LC-MS/MS determination of chloramphenicol in food of animal origin in Brazil; *ScientiaChromatographica*, 7(4):287-295.

20.SichilongoaK. F,Vallerie A,Muckoyab and Mathew M. N. **(2015)**, A Rapid and sensitive LC-MS/MS method for the determination of multi-class residues of antibiotics in chicken liver,*S. Afr. J. Chem.*, 68, 1–6.

21.Zhou J, Yao L, Yi L, Lanzhen C, Liming W. and Jing Z.**(2014)**, Floral classification of honey using liquid chromatography–diode array detection–tandem mass spectrometry and chemometric analysis,*Food Chemistry* 145:941–949.

22. WangS, **Yong** W, Jiahui L, Liya Z, Qilong C and Yiyang D.**(2014)**,Development of an indirect competitive assay-based aptasensor for highly sensitive detection of tetracycline residue in honey, *Biosensors and Bioelectronics* 57, 192-198.

23.MahmoudiR,Mirhasan M, Reza N,Sepideh K, Mohammad R. A. N. andKarim M.**(2014),** Detection of oxytetracycline residues in honey samples using ELISA and HPLC methods,*Pharmaceutical sciences*,19(4), 145-150.

24. **Fedorova G**, Nebesky V, randak T, Roman G. Simultaneous determination of 32 antibiotics in aquaculture products using LC-MS/MS, *Chemical Papers*, 68(1).

25.ZaiI. M, Khaliqur R,Arshad H. and Shafqatullah.**(2013)**, Detection and quantification of antibiotics residues in honey samples by chromatographic techniques,*Middle-East Journal of Scientific Research* 14 (5), 683-687

26.**AlechagaE,** Encarnacion M. and Galceran M. T. (2012),Ultra-high performance liquid chromatography-tandem mass spectrometry for the analysis of phenicol drugs and florfenicol-amine in foods ***Analyst,* 137,** 2486-2494.

27.**PerezaM. L. G**, Patricia P.N, Roberto R, Jose L. M. and Antonia G.**(2012)**, Comprehensive qualitative and quantitative determination of pesticides and veterinary drugs in honey using liquid chromatography–Orbitrap high resolution mass spectrometry,*Journal of Chromatography A*, 1248,130– 138.

28.Biea M, Rong L,Tingting C,Shouhui D,Hualin Z,Shuming Y.and Jing Q. (2012), Simultaneous determination of tetracycline antibiotics in beehives by liquid chromatography–triple quadrupole mass spectrometry, *Advances in Applied Science Research*, 3(1), 462-468.

29. CronlyM , Behan P, Foley B, Malone E , Martin S, Doyle M. and Regan L.**(2010)**, Rapid multi-class multi-residue method for the confirmation of chloramphenicol and eleven nitroimidazoles in milk and honey by liquid chromatography-tandem mass spectrometry (LC-MS), *Food Additives & Contaminants: Part A* 27(9), 1-20.

30.**Taokaenchan**N. and Supaporn S.**(2010)**,HPLC-Fluorescence detection method for quantitative determination of tetracycline antibiotic residues in honey,*NU Science Journal* 6(2), 147-155.

31. Vidal J. L.M., Luiz, M. M. A., Gonzalez R. R. and Frenich A. G. (2009) Multiclass analysis of antibiotic residues in honey by ultra-performance liquid chromatographytandem mass spectrometry, *J. Agri. Food Chem*. **57** (5), 1760–1767.

32. Pan C, ZhangH, ChenS,Xu Y. and Jiang S. **(2006)**,Determination of chloramphenicol residues in honey by monolithic column liquid chromatography–mass spectrometry after use of quechers clean-up,*ActaChromatographica*, 17, 320-327.

33.GallinaA.BenettiC,Giancasrlo B. and Franco M.**(2005)**, Antibiotic residues in honey: validation procedure, *APIACTA* 40, 45-49.

34. Bogialli S, Roberta C, Antonio Di C, Aldo L, Monica M and Manuela N.**(2005)**, Simple confirmatory assay for analyzing residues of aminoglycosideantibiotics in bovine milk: hot water extraction followed by liquidchromatography–tandem mass spectrometry; *Journal of Chromatography A*, 1067, 93–100.

35. Ortelli D, Edder P. and Corvi C.**(2004)**, Analysis of chloramphenicol residues in honey byliquid chromatography–tandem mass spectrometry, *Chromatographia*59(1), 61-64.

36. Pagliuca G., Gazzotti T., Serra G and Sabatini A.G.**(2002),** A scientific note on thedetermination of oxytetracycline residues in honey by high-performance liquidchromatography with UV detection, *Apidologie*,**33,** 583-584.